

Strange Isosinglet Versus Octet Scalar Axial Current

M. Kirchbach

Institut für Kernphysik, J. Gutenberg Universität, D-55099 Mainz, Germany

The flavor $SU(3)_f$ group symmetry of QCD is systematically considered as the heavy c quark limit of $SU(4)_f$. Within that scheme we argue that the $U(1)_A$ anomaly prevents Gell-Mann's choice for the realization of the $su(4)_f$ (and thereby the $su(3)_f$) algebra for the particular case of the neutral axial currents. Rather, Weyl's choice for the $SU(4)_f$ generators has to be used which leads to a neutral strong axial current having same structure (up to a constant factor) as the weak one.

PACS: 11.30.Hv, 11.40.Ex, 12.39.Jh

KEY words: flavor symmetry, hypercharge axial current, $U(1)_A$ anomaly

The $SU(3)_f$ flavor symmetry is one of the most important concepts of contemporary hadron physics. It has its roots in the empirical observation that the lowest mass pseudoscalar mesons and spin-1/2 baryons join to octets, while the lightest spin-3/2 baryons and the lightest vector mesons constitute a decuplet, and a nonet, respectively. From this point of view the existence of the first three fundamental flavor degrees of freedom of hadron matter, the u , d , and s quarks, has been concluded. After the discovery of the charmed (c) quark, the extension of $SU(3)_f$ to $SU(4)_f$ was considered despite of the apparent mismatch between the mass of the c quark on the one side, and the masses of the u , d , and s quarks, on the other side. Nonetheless, this extension is useful because the $SU(4)_f$ group represents a symmetry operation acting onto two complete quark generations and thus has common representations with the electroweak symmetry. Henceforth, the $SU(3)_f$ group which still preserved its importance as the relevant flavor symmetry of QCD, will be considered to emerge in the heavy c -quark limit of $SU(4)_f$.

The algebra of the four-flavor special unitary group is determined by the fifteen generators G^i satisfying the commutation relations

$$[G^i, G^j] = 2if_k^{ij} G^k, \quad i, j, k = 1, \dots, 15. \quad (1)$$

Note that the structure constants $f_k^{ij} = \text{tr}([G^i, G^j]G_k)/4i$ (they are completely antisymmetric in the indices i, j , and k) depend on the explicit realization of the generators G^i as linearly independent traceless Hermitian operators.

The most natural realization of an $su(n)$ Lie algebra is constructed within the (n^2-1) -dimensional basis defined in terms of the matrices E_{ik} as [1]

$$\begin{aligned} \langle m | E_{ik} | n \rangle &= \delta_{im} \delta_{mk}, \quad i, k = 1, \dots, n, \quad i \neq k, \\ E'_{tt} &= E_{tt} - E_{t+1, t+1}, \quad t = 1, \dots, n-1, \\ [E_{ik}, E_{lm}] &= \delta_{im} E_{lk} - \delta_{lk} E_{im}. \end{aligned} \quad (2)$$

In other words, E_{ik} is a matrix containing the unit at the intersection between the k th row and the i th column, while all the other matrix elements are equal to zero. For the diagonal matrices E'_{tt} one finds $(E'_{tt})_{tt} = 1$, and $(E'_{tt})_{t+1, t+1} = -1$, respectively. The so-called Weyl choice for the $SU(4)$ generators is now given by

$$\begin{aligned} G^r &= E_{lm} + E_{ml}, \quad r = 1, \dots, 6, \quad \text{for } l < m, m = 2, \dots, 4, \\ G^s &= E'_{k,k}, \quad s = 7, 8, 9, \quad \text{for } k = 1, 2, 3, \\ G^t &= i(E_{lm} - E_{ml}), \quad t = 10, \dots, 15, \quad \text{for } l < m, m = 2, \dots, 4. \end{aligned} \quad (3)$$

Note that any set of 15 matrices obtained as linearly independent combinations of the Weyl matrices defined above can be considered as a realization of the $su(4)_f$ algebra. For example, one can construct a new $su(4)_f$ algebra in keeping all the non-diagonal elements unchanged while replacing the diagonal elements E'_{11} , E'_{22} and E'_{33} by the respective matrices Λ^3 , Λ^8 , and Λ^{15} introduced as

$$\begin{aligned} \Lambda^3 &= E'_{11} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ \Lambda^8 &= \frac{1}{\sqrt{3}} (E'_{11} + 2E'_{22} + 2E'_{33}) = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix}, \quad \Lambda^{15} = \frac{1}{\sqrt{6}} (E'_{11} + 2E'_{22} - E'_{33}) = \frac{1}{\sqrt{6}} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -3 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \end{aligned} \quad (4)$$

This new set of $SU(4)_f$ generators is known from standard textbooks as Gell-Mann's choice for the $su(4)_f$ algebra (see Ref. [2]). The question which one of these two algebras is relevant for hadron systems can be answered only by comparison with suitable physical observables. We here show that, while both algebra choices appear equivalent for the vector currents, this is not the case for the axial vector currents. In the latter case, the $U(1)_A$ anomaly prevents the existence of an octet axial current and thereby Gell-Mann's choice for the $su(4)_f$ algebra. Our argumentation goes as follows.

The main physical manifestations of unitary symmetries are twofold. These are the particle multiplets, on the one side, and the Noether currents, on the other side. From the mathematical foundations of Lie groups it is well known, that while the dimensionality of the irreducible representations of the group under consideration does not depend on the concrete realization of its generators, the Noether currents do. To gain an insight into the choice for the diagonal $su(4)_f$ elements relevant for hadron processes one has, therefore, to analyze the flavor structure of the fundamental electromagnetic (vector) current, J_μ of the quarks. In doing so, the following decomposition of J_μ is found in the Weyl basis:

$$\begin{aligned} J_\mu &= \frac{2}{3}\bar{u}\gamma_\mu u - \frac{1}{3}\bar{d}\gamma_\mu d + \frac{2}{3}\bar{c}\gamma_\mu c - \frac{1}{3}\bar{s}\gamma_\mu s \\ &= J_\mu^{11} + J_\mu^{33} + J_\mu^B, \\ J_\mu^{11} &:= \bar{q}\gamma_\mu \frac{E'_{11}}{2} q, \quad J_\mu^{33} := \bar{q}\gamma_\mu \frac{E'_{33}}{2} q, \quad J_\mu^B := \frac{1}{2}\bar{q}\frac{\mathbb{1}_4}{3}\gamma_\mu q, \quad \bar{q} = (\bar{u}, \bar{d}, \bar{c}, \bar{s}). \end{aligned} \quad (5)$$

Here, the current J_μ^{11} is precisely the third component of the ordinary isospin current, while J_μ^B is half the baryon number current. Special attention has to be paid to the last current J_μ^{33} appearing diagonal in the fundamental quark quadruplet. This current is non-vanishing only for the heavy flavor c and s quarks from the second quark generation, and its diagonal matrix elements in turn equal $\frac{1}{2}C$, and $\frac{1}{2}S$, with C and S standing in turn for the respective charm and strangeness quantum numbers of the quarks.

From Eq. (5) follows that the fundamental vector quark current J_μ transforms according to the sum of the singlet representation $\{0\}$ with the 1st and 2nd quark generation doublets $\{2\}_{ud}$, and $\{2\}_{cs}$, respectively, rather than according to the fundamental quark quadruplet $\{4\}$, i.e. one finds J_μ to transform according to

$$\{4\} \rightarrow \{2\}_{ud} \oplus \{2\}_{cs} \oplus \{0\}. \quad (6)$$

This decomposition shows that the symmetry of the QCD Lagrangian in the neutral sector is in fact not the full $SU(4)_f$ group but rather its restriction to $SU(2)_{ud} \otimes SU(2)_{cs} \otimes U(1)_V$. Here the $SU(2)_{ud}$ and $SU(2)_{cs}$ groups act in turn onto the 1st and the 2nd quark generations. On the level of the algebra one has

$$su(4)_f \rightarrow su(2)_{ud} \oplus su(2)_{cs} \oplus u(1)_V. \quad (7)$$

As long as both J_μ^{33} and the baryon current are conserved, one can introduce their sum, here denoted by J_μ^Y , as a new conserved current

$$J_\mu^Y = J_\mu^{33} + J_\mu^B. \quad (8)$$

The total flavor charge, $Q^Y(t)$, of that current

$$Q^Y(t) = \int_0^\infty J_0^Y(t, \vec{x}) d^3\vec{x}, \quad (9)$$

will be a constant of motion and can be simultaneously diagonalized with the Hamiltonian of the hadron system. Its eigenvalues for the 1st and 2nd quark generations are given in turn by the following expressions,

$$Q^Y \begin{pmatrix} c \\ s \end{pmatrix} = \begin{pmatrix} C + \frac{1}{3} & 0 \\ 0 & S + \frac{1}{3} \end{pmatrix} \begin{pmatrix} c \\ s \end{pmatrix}, \quad Q^Y \begin{pmatrix} u \\ d \end{pmatrix} = \begin{pmatrix} \frac{1}{3} & 0 \\ 0 & \frac{1}{3} \end{pmatrix} \begin{pmatrix} u \\ d \end{pmatrix}, \quad (10)$$

and can be used to label the hadron states. From Eq. (5) follows that the total vector charge operator $\hat{Q}_V(t) = \int_0^\infty J_0(t, \vec{x}) d^3\vec{x}$ is related to the operators of isospin $\hat{t}_3 = \frac{1}{2}E'_{11}$, and hypercharge, $\hat{Y} = E'_{33} + \frac{1}{3}\mathbb{1}_4$, via

$$\hat{Q}_V = \hat{t}_3 + \frac{\hat{Y}}{2} = \frac{1}{2} \begin{pmatrix} 1 + \frac{1}{3} & 0 & 0 & 0 \\ 0 & -1 + \frac{1}{3} & 0 & 0 \\ 0 & 0 & C + \frac{1}{3} & 0 \\ 0 & 0 & 0 & S + \frac{1}{3} \end{pmatrix} \quad (11)$$

thus giving rise to the Gell-Mann–Nishijima relation. Eq. (11) clearly illustrates that four-flavor hypercharge can not be introduced on the $SU(4)_f$ level but rather requires the full $U(4)_f$ group.

Now in freezing the charm degree of freedom of the hypercharge current in Eq. (8), Gell-Mann’s octet vector current is reproduced in the truncated flavor space as

$$\frac{1}{2\sqrt{3}}J_\mu^8 = \lim_{m_c \rightarrow \Lambda_c} J_\mu^Y, \quad \Lambda_c \approx 1.5 \text{ GeV}. \quad (12)$$

Here Λ_c has been chosen to be sufficiently large in order to ensure negligible c quark effects on the 1 GeV mass scale, on the one side, but finite, in order to preserve the anomaly free character of the $SU(4)_f$ theory, on the other side [3]. Eq. (12) creates the impression, that for the particular case of three flavors, hypercharge may be introduced within the special group. In fact this is not correct as the essential ingredient of the hypercharge is the baryon number which, in being associated with the conservation of the $U(1)_V$ current, can not be defined within the $SU(3)_f$ framework. In other words, if Λ^8 is to enter the $SU(3)_f$ version of the Gell-Mann–Nishijima relation and be interpreted as the hypercharge generator, it has to be defined instead by Eq. (4) rather by

$$\frac{1}{\sqrt{3}}\Lambda^8 = \lim_{m_c \rightarrow \Lambda_c} \left(E'_{33} + \frac{1}{3} \mathbb{1}_4 \right). \quad (13)$$

Note that by means of Λ^8 from Eq. (4) the electric charges of the quarks would be obtained as an artificial combination of third projections of isospin (t_3) and U-spin ($t_3^U := E'_{22}/2$) quantum numbers according to $Q = t_3 + \frac{1}{3}(t_3 + 2t_3^U)$ rather than by the genuine Gell-Mann–Nishijima relation in Eq. (11). Until one is concerned with the anomaly free vector currents, however, Gell-Mann’s matrix Λ^8 can be viewed as an *effective* representation of hypercharge within the truncated flavor space. In such a case, using Gell-Mann’s choice for the $su(3)_f$ algebra doesn’t create any problems and is equivalent to Weyl’s choice. In the axial sector, however, Gell-Mann’s choice is of minor use as the axial transformation associated with the unit matrix from Eq. (13) will transport the problems of the $U(1)_A$ anomaly onto the octet axial current.

The observation that the octet matrix Λ^8 has no independent fundamental meaning as an element of the $su(4)_f$ algebra is further strongly supported by the internal flavor structure of the neutral vector mesons which are evidently Weyl bosons as their states transform in accordance to $SU(2)_{ud} \otimes SU(2)_{cs} \otimes U(1)_V$ representations:

$$\begin{aligned} |\rho^0\rangle &= \frac{1}{\sqrt{2}}(\bar{u} \bar{d})E'_{11} \begin{pmatrix} u \\ d \end{pmatrix}, \quad |\omega\rangle = \frac{1}{\sqrt{2}}(\bar{u} \bar{d})\mathbb{1}_2 \begin{pmatrix} u \\ d \end{pmatrix}, \\ |\phi\rangle &= \lim_{m_c \rightarrow \Lambda_c} (\bar{c} \bar{s})E'_{33} \begin{pmatrix} c \\ s \end{pmatrix}. \end{aligned} \quad (14)$$

The well known ‘ideal mixing’ within the vector meson nonet therefore corresponds to the rotation of Gell-Mann’s to Weyl’s basis. Note that the separation of the strange and non-strange quarkonia within the vector meson nonet is known as the Okubo-Zweig-Iizuka (OZI) rule and reflects the existence of gluon exchange between the quarks. The OZI rule is therefore an expression for favoring Weyl’s over Gell-Mann’s choice for $su(4)_f$.

Now one has to answer the question whether or not Gell-Mann’s choice for the elements of the $su(3)_f$ algebra will apply to the axial vector currents. To answer this question let us first recall that the generators of axial flavor transformation don’t constitute an algebra by themselves. They can mostly be considered as ingredients of group transformations when associating them to the vector flavor transformation to produce the respective left (L) and right (R) chiral rotations. For this reason, the axial currents *have to replicate in flavor space the structure of the vector currents*. In other words, the neutral chiral hadron currents will also transform according to the representation of

$$[su(2)_{ud}^{(L)} \oplus su(2)_{cs}^{(L)}] \oplus [su(2)_{ud}^{(R)} \oplus su(2)_{cs}^{(R)}] \oplus [u(1)_V \oplus u(1)_A]. \quad (15)$$

Within the context of Eq. (15), the only neutral axial current having a well defined chiral limit will be

$$J_{\mu,5} = \bar{q}\gamma_\mu\gamma_5\frac{E'_{11}}{2}q + \bar{q}\gamma_\mu\gamma_5\frac{E'_{33}}{2}q. \quad (16)$$

The decomposition of $J_{\mu,5}$ in the last equation reflects the exclusion of the anomalous $U(1)_A$ current [4] which can not be used any longer as a building block for the construction of an anomaly free octet axial current.

One remarkable feature of $J_{\mu,5}$ is that its structure is identical (up to the factor of $-1/2$) to that of the weak axial vector current. For this reason, the well established universality of the charged weak and strong axial vector currents underlying the current algebra can be extended to include the neutral ones. In the heavy c quark limit, the $J_{\mu,5}$ current decomposes into an isovector ($J_{\mu,5}^I$) and a purely strange $SU(2)_I$ isosinglet ($J_{\mu,5}^s$) component

$$J_{\mu,5}^I = \bar{q}\gamma_\mu\gamma_5\frac{E'_{11}}{2}q, \quad J_{\mu,5}^s = \lim_{m_c \rightarrow \Lambda_c} (\bar{q}\gamma_\mu\gamma_5\frac{E'_{33}}{2}q). \quad (17)$$

In contrast to the physical vector mesons ϕ and ω , their corresponding η and η' mesons from the pseudoscalar nonet are not Weyl bosons respecting the OZI rule. Within the 0^- nonet the $U(1)_A$ anomaly leads to a significant mixing of the Weyl states and thereby to a violation of the OZI rule [11]. For the particular case of quadratic mass formulae, the physical η meson turns out to emerge in the heavy c quark limit from the following Weyl states mixing:

$$|\eta\rangle = \lim_{m_c \rightarrow \Lambda_c} \cos\epsilon (\bar{c}c - \bar{s}s) - \sin\epsilon \frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d), \quad (18)$$

with $\epsilon \approx -45, 40^\circ$. This is an alternative way to express the tendency of the flavor structure of the η meson towards Gell-Mann-Okubo mass formulae predictions. In other words, while the wave function of the η meson can be roughly associated with the scalar of the octet, the ηNN vertex will be an isosinglet rather than an F-spin scalar because the η meson has to couple to the isosinglet nucleon current,

$$J_{\mu,5}^{s(N)} = \langle N | \frac{1}{2} \bar{s}\gamma_\mu\gamma_5 s | N \rangle = G_1^s \bar{\mathcal{U}}_N \gamma_\mu \gamma_5 \frac{\mathbb{1}}{2} \mathcal{U}_N, \quad G_1^s = \Delta s, \quad (19)$$

which can happen only via its strange quarkonium component. In assuming, for simplicity, the wave function of the η meson to be the pure scalar of the octet, its isosinglet axial current is now defined as

$$J_{\mu,5}^{s(\eta)} = \langle 0 | \frac{1}{2} \bar{u}\gamma_\mu\gamma_5 u - \frac{1}{2} \bar{d}\gamma_\mu\gamma_5 d - \frac{1}{2} \bar{s}\gamma_\mu\gamma_5 s | \frac{1}{\sqrt{6}}(\bar{u}u + \bar{d}d - 2\bar{s}s) \rangle = f_\eta i q_\mu, \quad f_\eta := \frac{1}{\sqrt{6}} \kappa_s(0^-) m_\eta. \quad (20)$$

Here $\kappa_s(0^-)$ denotes the dimensionless coupling of the strange quarkonium component of the pseudoscalar meson to the strange axial vector current, while f_η , m_η and q_μ are in turn the weak decay coupling, the mass and the momentum of the η meson. Furthermore, the couplings between $\bar{u}u$ and $\bar{d}d$ quarkonia and quark axial currents have been suggested to be diagonal in flavor and universal in strength, i.e. $\kappa_u(0^-) = \kappa_d(0^-)$, to recover isospin symmetry already at the tree level¹. With this in mind, the contact ηNN vertex takes the form

$$\begin{aligned} \mathcal{V}_{\eta NN} &= \frac{1}{f_\eta^2} J^{s(N)} \cdot J^{s(\eta)} \\ &= \frac{1}{f_\eta^2} \left(G_1^s \bar{\mathcal{U}}_N \gamma_\mu \gamma_5 \frac{\mathbb{1}}{2} \mathcal{U}_N \right) \cdot f_\eta i q_\mu. \end{aligned} \quad (21)$$

Here, the parameter of dimensionality $[mass^2]$ entering the current-current coupling has been chosen to equal the weak η decay coupling constant. This choice is convenient because of the universality of the isosinglet axial current and the related ansatz on the meson pole dominance. It allows one to express the coupling of neutral pseudoscalar mesons to arbitrary targets in terms of their weak decay couplings to the vacuum. The combination $\frac{G_1^s}{2f_\eta}$ is conventionally denoted by $\frac{f_{\eta NN}}{m_\eta}$ with $f_{\eta NN}$ being called the gradient ηN coupling constant. In inserting Eqs. (19) and (20) in the last expressions, one is led to

$$\begin{aligned} \frac{f_{\eta NN}}{m_\eta} &= \frac{G_1^s}{2f_\eta} = \sqrt{6} \frac{\Delta s}{2\kappa_s(0^-)m_\eta}, \\ f_{\eta NN} &= \sqrt{\frac{3}{2}} \frac{\Delta s}{\kappa_s(0^-)}. \end{aligned} \quad (22)$$

This means that at tree level the contact gradient ηN coupling appears proportional to the fraction of proton helicity carried by the strange quark sea, rather than to the octet axial vector coupling $g_A^{(8)} = \frac{1}{\sqrt{3}}(\Delta u + \Delta d - 2\Delta s)$, a result

¹ The empirically observed closeness of the π and η weak decay constants, ($f_\eta \approx 1.1 f_\pi$) does not necessarily imply $\kappa_u(0^-) \approx \kappa_s(0^-)$ at the tree level.

already conjectured in a previous work [5]. This might be one of the main reasons for which a strong suppression of the gradient ηN coupling has constantly been found over the years by various data analyses regarding the η photoproduction off proton at threshold [6], the $\bar{p}p$ collisions [7], as well as the nucleon-nucleon (NN) and nucleon-hyperon (NY) phase shifts [8].

The essence of the present study is that to determine the ηN coupling one has to exploit the neutral current in Eq. (16) which respects the OZI rule and can be considered formally to emerge from an ideal mixing between Gell-Mann's octet and flavor singlet axial currents. That $\eta_8 - \eta_0$ mixing will lead to a reduction of the nucleon matrix element of the octet axial current was earlier considered, for example, in Ref. [9]. There, however, the reduction reported was not as big as the one in Eq. (22).

In view of the small ηN contact coupling, loop corrections acquire importance. The effect of one-loop corrections on the suppression of the nucleon matrix element of the octet axial vector current has been considered in Ref. [10] within the framework of chiral perturbation theory. There, the strong dependence of the result on the nucleon $-\Delta$ mass splitting was revealed and the necessity for higher order corrections discussed. In Ref. [5] vertex renormalization effects were considered in terms of nucleon and meson degrees of freedom rather than on the quark level. There, the significance of triangular corrections of the type $a_0(980)\pi N$ for the enhancement of the $(\bar{s}s)NN$ vertex was revealed. These provide the coupling of the strange quarkonium component of the η meson to the non-strange meson cloud surrounding the nucleon via the violation of the OZI rule and lead to an effective ηN coupling, less than half the value predicted by the constituent quark model, in line with data.

To conclude, it was shown that the only isoscalar axial vector current having a really anomaly free chiral limit is $J_{\mu,5}$ in Eq. (16) which, in being a Weyl current, automatically respects the OZI rule. Unfortunately, due to $U(1)_A$ anomaly effects, the internal flavor structure of the η meson is not in accord with $J_{\mu,5}^s$. For these reasons, the physical η meson can mostly be considered as an approximate 'strange' Goldstone boson. As the $U(1)_A$ anomaly was shown to prevent the construction of an octet axial current with a well defined chiral limit, the interpretation of the η meson as the octet Goldstone boson has to be dropped despite the presence of $(\bar{u}u + \bar{d}d)$ components in its wave function.

Helpful comments by Martin Reuter and Andreas Wirzba are kindly appreciated. Work supported by the Deutsche Forschungsgemeinschaft (SFB 201).

-
- [1] J. F. Cornwell, *Group Theory in Physics* (Techniques in Physics, 1984).
 - [2] W. Greiner and B. Müller, *Quantenmechanik, Symmetrien* (Harri Deutsch, Frankfurt M., 1990), p. 429.
 - [3] J. Ambjoern, J. Greensite, and C. Peterson, Nucl. Phys. **B221**, 381 (1983).
 - [4] G. 't Hooft, Phys. Rev. **D14**, 3432 (1976).
 - [5] M. Kirchbach and H.-J. Weber, Comm. Nucl. Part. Phys. **22**, 171 (1998).
M. Kirchbach and L. Tiator, Nucl. Phys. **A604**, 385 (1996).
S. Neumeier and M. Kirchbach, MKPH-T-98-13.
 - [6] B. Krusche et al., Phys. Rev. Lett. **74**, 3736 (1995);
L. Tiator, C. Bennhold and S. Kamalov, Nucl. Phys. **A580**, 455 (1994).
 - [7] W. Grein and P. Kroll, Nucl. Phys. **A338**, 332 (1980).
 - [8] A. Reuber, K. Holinde, and J. Speth, Nucl. Phys. **A570**, 541 (1991).
 - [9] G. Veneziano, Mod. Phys. Lett. **A4** (1989) 1605.
 - [10] M. J. Savage and J. Walden, Phys. Rev. **D55**, 5376 (1997).
 - [11] S. Forte and E. Shuryak, Nucl. Phys. **B367**, 153 (1991).